

S.S. Gershtein, A.A. Logunov, and M.A. Mestvirishvili

UPPER LIMIT ON THE GRAVITON MASS

The problem of the existence of nonzero invariant mass of the graviton can have the fundamental significance. The estimate for the upper bound on the graviton mass ($m_g < 2 \cdot 10^{-62}$) has been derived in Ref. [1], where the authors used the data on the existence of the gravitational coupling between the galaxy clusters, which is not cut off by the Yukawa potential at least up to distances ~ 500 Kpc. In this paper we will present the estimates for upper limit on the graviton mass basing on the observable parameters of the Universe expansion. The fact that in this case typical distances are 3-4 orders of magnitude larger than those between gravitationally bound galaxy clusters allows one to strengthen the estimates on the upper limit on the graviton mass by few orders of magnitude, respectively.

It should be noted that introduction of the nonzero invariant mass of the graviton requires to go beyond the General Theory of Relativity (GTR). This can be done naturally by using the notions of the gravitational field in the Minkowsky space [2,3]. In Ref. [3] the complete energy-momentum tensor $t^{\mu\nu}$ (including the gravitational field), which is conserved in the Minkowsky space, is considered as a source of the gravitational field described by symmetric tensor $\Phi^{\mu\nu}$. In arbitrary fixed (not necessarily inertial) frame of the Minkowsky space with metric tensor $\gamma_{\mu\nu}$ the equations for the density of the gravitational field $\tilde{\Phi}^{\mu\nu}$ can be written analogously to the Maxwell equations and Lorentz condition for the electromagnetic field as follows:

$$(\gamma^{\alpha\beta} D_\alpha D_\beta + m_g^2) \tilde{\Phi}^{\mu\nu} = 16\pi \tilde{t}^{\mu\nu}, \quad (1)$$

$$D_\nu \tilde{\Phi}^{\mu\nu} = 0, \quad (2)$$

where D_α is the covariant derivative in the Minkowsky space, m_g is the graviton mass ($\hbar = c = G = 1$), and $\tilde{\Phi}^{\mu\nu}, \tilde{t}^{\mu\nu}$ are the densities of tensors:

$$\tilde{\Phi}^{\mu\nu} = \sqrt{-\gamma} \Phi^{\mu\nu}, \quad \tilde{t}^{\mu\nu} = \sqrt{-\gamma} t^{\mu\nu}, \quad \gamma = \det(\gamma_{\mu\nu}) = \det(\tilde{\gamma}^{\mu\nu}). \quad (3)$$

Condition (2) singles out polarization states with spin values of 2 and 0 and provides the conservation of the density of the energy-momentum tensor

$D_\mu \tilde{t}^{\mu\nu} = 0$ in Eq. (1). The density of the energy-momentum tensor is defined, following Hilbert, by Euler's variation with $\gamma_{\mu\nu}$ metric of the Lagrangian density of the system

$$\tilde{L} = \tilde{L}_g(\gamma_{\mu\nu}, \Phi_{\mu\nu}) + \tilde{L}_M(\gamma_{\mu\nu}, \Phi_{\mu\nu}, \Phi_A), \quad (4)$$

where $\tilde{L}_g$ is the density of the gravitational field Lagrangian, and $\tilde{L}_M$ corresponds to the density of Lagrangian of the matter described by the Φ_A fields.

$$\tilde{t}^{\mu\nu} = -2 \frac{\delta \tilde{L}}{\delta \gamma_{\mu\nu}}, \quad (5)$$

where Euler's variation is

$$\frac{\delta \tilde{L}}{\delta \gamma_{\mu\nu}} = \frac{\delta \tilde{L}}{\delta \gamma_{\mu\nu}} - \partial_\sigma \left(\frac{\partial \tilde{L}}{\partial \gamma_{\mu\nu,\sigma}} \right); \gamma_{\mu\nu,\sigma} = \frac{\partial \gamma_{\mu\nu}}{\partial x^\sigma}. \quad (6)$$

On can derive equations for the gravitational field and matter fields from the least action principle

$$\frac{\delta \tilde{L}}{\delta \tilde{\Phi}^{\mu\nu}} = 0, \quad \frac{\delta \tilde{L}}{\delta \Phi_A} = 0. \quad (7)$$

To get the form of (1) and (2) for these equations it is necessary to have the density of the gravitational field $\tilde{\Phi}^{\mu\nu}$ coming into the density of the matter Lagrangian $\tilde{L}_M$ in combination with the density of the metric tensor $\tilde{\gamma}^{\mu\nu}$:

$$\tilde{g}^{\mu\nu} = \tilde{\gamma}^{\mu\nu} + \tilde{\Phi}^{\mu\nu}, \quad \tilde{g}^{\mu\nu} = \sqrt{-g} g^{\mu\nu}, \quad g = \det(g_{\mu\nu}) = \det(\tilde{g}^{\mu\nu}), \quad (8)$$

i.e. $\tilde{L}_M(\tilde{g}^{\mu\nu}, \Phi_A)$. It means that the motion of the matter subjected to the gravitational field looks like as it could take place in the Riemann space with the metric $g_{\mu\nu}$. The Lagrangian density resulting to Eqs. (1) and (2) has the form

$$\tilde{L} = \tilde{L}_g + \tilde{L}_M(\tilde{g}^{\mu\nu}, \Phi_A), \quad (9)$$

$$\tilde{L}_g = \frac{1}{16\pi} \tilde{g}^{\mu\nu} (G_{\mu\nu}^\lambda G_{\lambda\sigma}^\sigma - G_{\mu\sigma}^\lambda G_{\nu\lambda}^\sigma) - \frac{m^2}{16\pi} \left(\frac{1}{2} \gamma_{\mu\nu} \tilde{g}^{\mu\nu} - \sqrt{-g} - \sqrt{-\gamma} \right), \quad (10)$$

where the $G_{\mu\nu}^\lambda$ values are the components of the tensor

$$G_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (D_\mu g_{\nu\sigma} + D_\nu g_{\mu\sigma} - D_\sigma g_{\mu\nu}), \quad (11)$$

and due to this fact the $\tilde{L}_g$ value behaves as the density of the scalar under any coordinate transformations. Using (9) and (10) and taking into account (7) one can write the equations for the gravitational field in the form of [3]

$$\left(R_\nu^\mu - \frac{1}{2}\delta_\nu^\mu R\right) + \frac{m_g^2}{2}(\delta_\nu^\mu + g^{\mu\alpha}\gamma_{\alpha\nu} - \frac{1}{2}\delta_\nu^\mu g^{\alpha\beta}\gamma_{\alpha\beta}) = 8\pi T_\nu^\mu \quad (12)$$

$$D_\nu \tilde{g}^{\mu\nu} = 0, \quad (13)$$

where T_ν^μ is the matter energy-momentum tensor in the Riemann space.

From these equations one gets the equations for the matter

$$\nabla_\nu \tilde{T}^{\mu\nu} = 0, \quad \tilde{T}^{\mu\nu} = -2 \frac{\delta \tilde{L}_M}{\delta g_{\mu\nu}}, \quad (14)$$

where ∇_ν is the covariant derivative in the effective Riemann space. Eqs. (12) and (13) are covariant with respect to any coordinate transformations and form-invariant under Lorentz's transformations.

Writing down the interval of the effective Riemann space for the homogeneous and isotropic Universe in the form of

$$ds^2 = U(t)dt^2 - V(t) \left[\frac{dr^2}{1 - kr^2} + r^2(d\Theta^2 + \sin^2 \Theta d\Phi^2) \right], \quad (15)$$

(where $k = 1, -1, 0$ for the closed, hyperbolic and “flat” Universe), one gets from Eqs. (13)

$$\frac{\partial}{\partial t} \sqrt{\frac{V^3}{U}} = 0, \quad \dots V = aU^{1/3}, a = \text{const.} \quad (16)$$

$$\frac{\partial}{\partial r} [r^2(1 - kr^2)^{1/2}] - 2r(1 - kr^2)^{-1/2} = 0, \quad (17)$$

Eq. (17) is valid only for $k = 0$. Thus, the Universe can be only “flat” (i.e. its space geometry is Euclidian). Using the proper time $d\tau = U^{1/2}dt$ and denoting $R^2 = U^{1/3}$ one can write interval (15) in the form

$$ds^2 = d\tau^2 - aR^2(\tau)(dx^2 + dy^2 + dz^2). \quad (18)$$

In this case Eqs. (12) in inertial frame take the form¹

$$\left(\frac{1}{R} \frac{dR}{d\tau}\right)^2 = \frac{8\pi G}{3} \rho - \frac{\omega}{R^6} \left(1 - \frac{3R^4}{a} + 2R^6\right), \quad (19)$$

$$\frac{1}{R} \cdot \frac{d^2 R}{d\tau^2} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2}\right) - 2\omega \left(1 - \frac{1}{R^6}\right), \quad (20)$$

where

$$\omega = \frac{1}{12} \left(\frac{m_g c^2}{\hbar}\right)^2. \quad (21)$$

It follows from Eq. (19) in the region $R \gg 1$ that the density of the matter in the Universe is equal to

$$\rho(\tau) = \rho_c(\tau) + \frac{1}{16\pi G} \left(\frac{m_g c^2}{\hbar}\right)^2, \quad (22)$$

where $\rho_c(\tau)$ is the critical density determined by the Hubble “constant”:

$$\rho_c = \frac{3H^2(\tau)}{8\pi G}, \quad H(\tau) = \frac{1}{R} \cdot \frac{dR}{d\tau}. \quad (23)$$

This conclusion inevitably requires the existence of the “dark” matter that agrees with current observations.

From (19) and (20) one can get the expression for the Universe deceleration parameter $q(\tau)$. At the present stage of the nonrelativistic matter dominance ($p = 0$)

$$q = -\frac{\ddot{R}}{R} \cdot \frac{1}{H^2} = \frac{1}{2} + \frac{1}{4H^2} \left(\frac{m_g c^2}{\hbar}\right)^2. \quad (24)$$

¹Note, that the metric of the Minkowsky space $\gamma_{\mu\nu}$ comes into Eq. (12). Due to this fact the Minkowsky space becomes observable, and the casualty principle for the gravitational field in the effective Riemann space should be fulfilled: the motion of the matter subjected to the gravitational field should not leave the light-cone limits in the Minkowsky space. This condition can be formulated in the form of $g_{\mu\nu} V^\mu V^\nu \leq 0$ for any isotropic vector V^μ on the light cone $\gamma_{\mu\nu} V^\mu V^\nu = 0$. This condition being applied to interval (15) taking into account (16) and (17) leads to $R^2(R^4 - a) \leq 0$. Thus the constant “ a ” has the notion of the fourth power of maximal value of the scale factor: $a = R_{\max}^4$, and to describe the existing Universe one should have $a \gg 1$.

Relation (24) gives the principal possibility to determine the graviton mass from two other observables, H and q . The sensitivity of q to the graviton mass is due to the fact that a small value $\frac{1}{\lambda_g} = \frac{m_g c}{\hbar}$ comes into (24) multiplied by a large value $\left(\frac{c}{H}\right) = 9.25 \cdot 10^{27} \cdot h^{-1}$ cm, which is the Hubble radius of the Universe. Though the q value has not been measured with high accuracy, its possible values do not exceed few units ($q \leq 5$, see. [4]). This allows one to estimate from (24) the graviton mass

$$m_g \leq 1.7 \cdot 10^{-65} \cdot h \text{ (g.)}, \text{ where } 0.4 \leq h \leq 1 \quad (25)$$

$$\frac{\hbar}{m_g c} > 0.2 \cdot \frac{c}{H} = 2 \cdot 10^{27} \cdot h^{-1} \text{ (cm.)}. \quad (26)$$

Despite of the smallness of the upper limit in (25), nonzero graviton mass can have principal influence on the character of the Universe evolution. One can see from Eq. (19) that for $R \rightarrow 0$ the negative term $\frac{\omega}{R^6}$ in the right-hand side of the equation grows in absolute value faster than the matter density ($\rho \sim \frac{1}{R^4}$ for radiatively dominant stage). Therefore, from the condition of the nonnegative left-hand side of (19) it follows that the expansion should begin from some minimal value $R_{\min}$, which corresponds to $\frac{dR}{d\tau} = 0$. From other side, the expansion should stop at $R \gg 1$, when density (22) reaches its minimal value $\rho_{\min} = \frac{1}{16\pi G} \left(\frac{m_g c^2}{\hbar}\right)^2$, and after that the expansion is replaced by the compression process up to $R_{\min}$. So, nonzero graviton mass eliminates the cosmological singularity and leads to cyclic character of the Universe evolution. Such a character of the Universe evolution seemed to be promising for a number of authors (see, for example, [5]). The time of the Universe expansion from the maximal density to the minimal one is determined mostly by the stage of the nonrelativistic matter dominance, and it is equal to [6]

$$\tau_{\max} \simeq \sqrt{\frac{2}{3}} \cdot \frac{\pi \hbar}{m_g c^2}. \quad (27)$$

Accepting the value of $(10 - 15) \cdot 10^9$ years for the Universe age and using $\tau_{\max} \geq 20 \cdot 10^9$ years, on gets more strict limit on the graviton mass

$$m_g \leq 4.5 \cdot 10^{-66} \text{ (g.)}. \quad (28)$$

Equations (12) and (13) explain all known gravitational effects in the Solar system, which are attributed to the noninertial frame. It is well known

that the introduction of the graviton mass in the linear tensor theory is accompanied by the difficulty: the presence of “ghosts”. However, as it was shown in Ref. [7], this difficulty is eliminated in the framework of the nonlinear tensor theory described by Eqs. (12) and (13) under condition that gravitons spread in the effective Riemann space, rather than Minkowsky’s one (as it takes place in the linear theory). If this circumstance is sequently taken into consideration, one gets a positively determined flux of the gravitational energy when calculating the intensity.

References

- [1] A.S.Goldhaber and M.M.Nieto. Phys.Rev.D9, 1119 (1974).
- [2] See. “Feynman Lectures on Graviton” Addison-Werley Publishing Company (1995);
W.Thirring. Ann. of Phys. 16 (1961)96;
V.I.Ogievetsky and I.V.Polubarinov. Ann.of Phys. 35 (1965)167 and refs. therein.
- [3] A.A.Logunov. Uspekhi Fiz. Nauk. 165, 187 (1995). Physics-Uspekhi 38(2), 179 (1995).
A.A., M.A.Mestvirishvili. Preprint IHEP 95-95 (1995).
A.A.Logunov. “Relativistic Theory of Gravity and the Mach Principle”. Dubna 1997.
- [4] E.W.Kolb and M.S.Turner. “The Early Universe”. See. Fig.1.2, p.5 with reference to J.Kristian, A.Sandage and J.Westphal, Ap.J.221, 383 (1978).
- [5] A.D.Sakharov. Scientific papers. Moscow, 1995, p.414; ZhETF 83, 1233 (1982).
- [6] S.S.Gershtein, A.A.Logunov, and M.A.Mestvirishvili. Preprint IHEP 97-36 (1997).
- [7] Yu.M.Loskutov. Vestnik MSU, 1991, phys. ser. 32, 4, p.49;
Yu.M.Loskutov. Proc. of the VI-th Marsel Grosman Meeting on Gen.Rel. Part B. Kyoto, Japan, p.1658 (1991).